

Allfarm Piperazine – wormer for poultry.

Allfarm Piperazine is used for the treatment of round worms (*Ascaridia* spp.) – one of the most prevalent and important intestinal worms of poultry. It has a wide safety margin and is suitable for all chickens, turkeys, ducks and pigeons. Allfarm Piperazine comes as a water-soluble powder and should be administered to birds in their drinking water.

APVMA Approval Number: 45124.

When should you use Allfarm Piperazine?

What are the symptoms of round worm infection?

Round worms may cause a wide range of general symptoms, including diarrhoea, depression, decreased appetite, drooping wings, ruffled feathers and lethargy. These symptoms are shared by a number of poultry diseases, including coccidiosis, and faecal testing is required for a definitive diagnosis.

Parasite Diagnostic Services performs combined worm egg and coccidiosis testing of poultry flocks to help you decide if your birds require treatment, or if previous treatment has been effective. See their website for more information, including pricing and contact details: <https://www.parasite.com.au/>

Active Ingredient:

Piperazine dihydrochloride (530 mg/g).

Directions:

Allfarm Piperazine is administered as a single treatment. If the round worm infestation is severe we recommend the treatment be repeated 17 to 21 days later.

- Determine the required amount of Allfarm Piperazine powder: use 1 g Piperazine powder per 5 kg of bird bodyweight (200 mg/kg). A 1 kg pack of Allfarm Piperazine treats 2500 birds with a bodyweight of 2 kg.
- Determine the required volume of drinking water: this is the amount that birds will consume within a 6-8 hour period. All treated water should be discarded after 8 hours.
- Premix the Allfarm Piperazine powder in a smaller volume of drinking water, dissolve completely and add to the medication tank or water supply, mixing thoroughly.
- Monitor the medicated water in case it is consumed quickly. If so, return birds to fresh drinking water.
- After 6-8 hours discard any remaining medicated water and return the birds to fresh drinking water.

Withholding period:

There is **no withholding period** for the use of Piperazine in poultry.

Storage:

Store in a cool, dry place ($\leq 30^{\circ}\text{C}$).

Shelf life:

Check the packaging for the expiry date.

First Aid:

If poisoning occurs contact a doctor, or call the Poisons Information Centre on 13 11 26 (Australia).

Ordering:

Allfarm products may be ordered from our website: <https://www.allfarm.com.au>
or by telephone: (03) 5979 4488.

Product	Code	Barcode
Allfarm Piperazine, 100 g	W00109W	9341382000587
Allfarm Piperazine, 1 kg	W00110W	9341382000242
Allfarm Piperazine, 5 kg	W00111W	9341382000259
Allfarm Piperazine, 15 kg	W00113W	9341382000266